

Hypercapnia Attenuates Hypoxic Pulmonary Hypertension by Inhibiting Lung Radical Injury

M. CHOVANEC^{1,4,5}, J. NOVOTNÁ^{3,4}, J. WILHELM^{3,4}, V. HAMPL^{1,4}, M. VÍZEK^{2,4}, J. HERGET^{1,4}

¹Department of Physiology and ²Department of Pathophysiology, ³Department of Biochemistry, Second Medical School, Charles University, Prague, ⁴Center for Cardiovascular Research, Prague, ⁵Cardiology Department, Na Homolce Hospital, Prague, Czech Republic

Received October 9, 2009

Accepted October 30, 2009

Summary

Chronic lung hypoxia results in hypoxic pulmonary hypertension. Concomitant chronic hypercapnia partly inhibits the effect of hypoxia on pulmonary vasculature. Adult male rats exposed to 3 weeks hypoxia ($F_{iO_2}=0.1$) combined with hypercapnia ($F_{iCO_2}=0.04-0.05$) had lower pulmonary arterial blood pressure, increased weight of the right heart ventricle, and less pronounced structural remodeling of the peripheral pulmonary arteries compared with rats exposed only to chronic hypoxia ($F_{iO_2}=0.1$). According to our hypothesis, hypoxic pulmonary hypertension is triggered by hypoxic injury to the walls of the peripheral pulmonary arteries. Hypercapnia inhibits release of both oxygen radicals and nitric oxide at the beginning of exposure to the hypoxic environment. The plasma concentration of nitrotyrosine, the marker of peroxynitrite activity, is lower in hypoxic rats exposed to hypercapnia than in those exposed to hypoxia alone. Hypercapnia blunts hypoxia-induced collagenolysis in the walls of prealveolar pulmonary arteries. We conclude that hypercapnia inhibits the development of hypoxic pulmonary hypertension by the inhibition of radical injury to the walls of peripheral pulmonary arteries.

Key words

Chronic hypoxia • Chronic hypercapnia • Pulmonary hypertension • Hypoxic lung injury

Corresponding author

Milan Chovanec, Department of Physiology, Second Medical School, Charles University, Prague, Plzeňská 221, 150 00 Prague 5, Czech Republic. E-mail: milan.chovanec@lfmotol.cuni.cz

Introduction

Chronic hypoxia causes hypoxic pulmonary hypertension (HPH) and structural remodeling of pulmonary blood vessels. Chronic lung hypoxia typically results from respiratory insufficiency or from breathing gas with a low oxygen concentration. The major difference between these two alternatives is in lung concentration of carbon dioxide. Whereas respiratory insufficiency in lung disease is most often characterized by hypercapnia and respiratory acidosis, oxygen deprivation in high mountains results in hypocapnia and alkalosis. In lung disease the onset of hypercapnia is an adverse prognostic event. Severity of cor pulmonale, however, was statistically linked with measurements of airway obstruction but not with hypercapnia (Wilkinson *et al.* 1988). In animal experiments, hypercapnia partially inhibited the development of HPH (Ooi *et al.* 2000, Herget *et al.* 2001, Herget *et al.* 2002, Howell *et al.* 2004, Kantores *et al.* 2006). In studies on high altitude dwellers (Canepa *et al.* 1956, Hultgren and Grover 1968) and patients with chronic obstructive pulmonary disease (COPD) (Lapp *et al.* 1971, Astin 1972), where both groups have similar levels of oxygen saturation of hemoglobin (~80%), high altitude residents have P_{aCO_2} ~31 torr and COPD patients ~46 torr. Hypercapnic sea level patients had less severe lung hypertension (mean pulmonary blood pressure 25 mm Hg) than mountain residents (35 mm Hg).

HPH results from hypoxia-induced radical tissue injury to the walls of peripheral pulmonary arterioles (Hampl and Herget 2000, Herget *et al.* 2000). The

Table 1. Body weight (BW), pulmonary artery pressure (PAP), cardiac output (CO), RV/LV+S ratio, DL ratio and plasma nitrotyrosine concentration in rats after 3 weeks exposure to ambient air (N), hypoxia (H) and hypoxia with hypercapnia (H + CO₂).

Group	BW (g)	PAP (mm Hg)	CO (ml.min ⁻¹)	RV/LV+S	DL (%)	Nitrotyrosine (μM)
N	415±8*	15.6±0.9*	38.4±4.2	0.252±0.012*	18±2*	6.9±0.6*
H + CO ₂	214±6	22.9±0.9 ⁺	38.0±3.3	0.349±0.016 ⁺	26±4 ⁺	11.8±1.4 ⁺
H	243±8	30.0±1.9	27.6±2.6	0.539±0.021	38±3	17.7±2.7

PAP – pulmonary arterial mean blood pressure, CO – cardiac output, RV/LV+S – ratio of right ventricular heart weight to the sum of weights of left heart ventricle and septum, DL – ratio of hypertrophied (double laminated) peripheral pulmonary arteries and total number of peripheral pulmonary arteries on section of both lungs. * P<0.05 between groups N and H + CO₂, ⁺ P<0.05 between groups H and H + CO₂.

initiation of HPH is characterized by the increase in production of NO and oxygen radicals (reactive oxygen species, ROS) (Lachmanová *et al.* 2005, Hampl *et al.* 2006). We hypothesize that hypercapnia restricts the HPH by inhibition of radical damage in the walls of the prealveolar pulmonary vessels. In this study on chronically hypoxic and chronically hypoxic and hypercapnic rats we characterized the developed pulmonary hypertension and analyzed the relation of hypercapnia to hypoxia-induced radical injury of peripheral pulmonary arteries.

Methods

Exposure to chronic hypoxia and hypercapnia

Experiments on three groups of adult male Wistar rats were performed in accordance with the European Community and NIH guidelines for using experimental animals. All procedures were approved by our institution's Animal Studies Committee.

The first group (group H) was exposed to chronic hypoxia in an isobaric hypoxic chamber (F_{iO₂}=0.1). CO₂ was completely reabsorbed in closed circuit by KOH and soda lime (Hampl and Herget 1990). In the second group (H + CO₂), rats were exposed to hypoxia (F_{iO₂}=0.1) and hypercapnia (F_{iCO₂}=0.04-0.05). The increase in carbon dioxide concentration was achieved by bypassing the KOH absorber. The concentration of CO₂ was continuously monitored. The third group of rats (N) lived in atmospheric air. Rats were exposed to hypoxia or to hypoxia and hypercapnia for 3 weeks (study A) or 4 days (study B).

Study A was designed to investigate the effect of hypercapnia on the development of hypoxic pulmonary hypertension. Study B was focused on the effects of

hypercapnia on ROS and NO production in early hypoxia-induced pulmonary vascular injury and on resulting metabolic changes in the walls of the peripheral pulmonary arteries.

After 3 weeks of exposure (study A) the rats were anesthetized with thiopental (30 mg/kg b.w. i.p.). In closed chest rats we measured the mean pulmonary arterial blood pressure (PAP) (Herget and Paleček 1972) and, after thoracotomy, the cardiac output (CO) by ultrasonic flow probe (Hampl *et al.* 2003). Hematocrit, plasma nitrotyrosine concentration (Herget *et al.* 2000) and concentration of lipofuscin-like pigments (lipid peroxidation endproducts) in erythrocyte membranes (Wilhelm and Herget 1999) were measured in samples of arterial blood. The heart was dissected and weighed in parts (Herget *et al.* 1978). The presence of chronic hypoxia-induced structural remodeling of peripheral pulmonary arteries was assessed by a quantitative histological method (Herget *et al.* 1978).

In rats exposed to 4 days hypoxia (study B) we measured the amount of exhaled NO in awake rats. After thiopental anesthesia, plasma concentration of NO plus its oxidation products (NO_x) and nitrotyrosine concentration were measured by methods described elsewhere (Hampl *et al.* 2006). Following thoracotomy the lungs were exposed, cooled with ice and peripheral pulmonary arteries were dissected. From 20-25 vascular samples obtained from individual rats we isolated collagen proteins by limited pepsin digestion and individual collagen fractions were analyzed by gel SDS electrophoresis (Novotná and Herget 1998).

Statistical analysis

Results were evaluated by ANOVA with Fischer *post-hoc* test. Values of p<0.05 were considered

significant. The results are presented as means \pm SEM.

Results

Study A – 3 weeks of exposure

Ten rats were included in each experimental group. All rats in groups N and H survived 3 weeks of experiment, one rat of the group H + CO₂ died after one week of exposure. The groups H and H + CO₂ had significantly lower body weight than normoxic controls N. The presence of high CO₂ concentration in inspired air significantly inhibited the development of HPH. Rats of the H + CO₂ group had significantly lower mean pulmonary arterial blood pressure, decreased right to left heart weight index and less muscularized peripheral pulmonary arteries than hypoxic rats with no hypercapnia (group H). Cardiac output did not differ between the groups. Hypercapnia, however, did not fully prevent the HPH and all indices of HPH were still significantly different from normoxic controls (Table 1). The values of hematocrit in groups exposed to hypoxia did not differ (H 74 \pm 2 %, H+CO₂ 73 \pm 3 %) and were significantly higher than in normoxic controls (N 54 \pm 3 %, P<0.001). The concentration of lipofuscin-like fluorescent pigments was significantly lower in the group H + CO₂ than in the group H (Fig. 1) and did not differ from normoxic controls.

Study B – 4 days of exposure

Four days exposure to hypoxia increased significantly the amount of NO in exhaled air. This increase was significantly lower in the group exposed to hypoxia and hypercapnia. Similarly the increase in plasma concentration of NO_x and serum concentration of nitrotyrosine was significantly lower in group H + CO₂ than in group H (Fig. 2).

Exposure to hypoxia induced collagenolysis in the walls of peripheral pulmonary arteries which resulted in the presence of typical low molecular weight collagen fragments (Novotná and Herget 1998). This hypoxia-induced collagen cleavage was less prominent in the group H + CO₂ than in the group H (Figs 3 and 4). Analysis of collagenous proteins was made in 7 rats of groups H and H + CO₂ and in 6 rats of group N.

Discussion

The main finding of this study is that hypercapnia inhibits the development of HPH by interaction with radical tissue injury in the walls of




Fig. 1. Concentration of lipofuscin-like fluorescent pigments after 3 weeks exposure to ambient air, hypoxia and hypoxia with hypercapnia. H – group exposed to isobaric hypoxia ($F_{iO_2}=0.1$), H + CO₂ – group exposed to isobaric hypoxia ($F_{iO_2}=0.1$) and hypercapnia ($F_{iCO_2}=0.04-0.05$), N – normoxic controls.

prealveolar pulmonary vessels. The finding that HPH is attenuated by an increase in carbon dioxide concentration is in agreement with previously published studies (Ooi *et al.* 2000, Herget *et al.* 2001, Herget *et al.* 2002). The rats exposed to combined chronic hypoxia and hypercapnia had significantly lower pulmonary arterial blood pressure, significantly lower increase in right heart ventricle weight and less pronounced structural remodeling of the walls of peripheral pulmonary arteries. In acute experiments, hypercapnia added during hypoxic pulmonary vasoconstriction caused further constriction or dilatation of lung vessels, depending on the initial vascular tonus and species of experimental animal (Emery *et al.* 1977). Ventilatory consequences of hypercapnia may alter the PaO₂ for given FiO₂ in short-term exposures (Pepelko 1970). We were not able to measure PaO₂ in our rats acclimatized to chronic exposure. However, the absence of differences in hematocrit values found in H and H + CO₂ groups suggests that the presence of an important difference in the level of hypoxemia is unlikely.

The present study indicates that the probable mechanism whereby hypercapnia inhibits HPH *via* the inhibitory effect of carbon dioxide on tissue hypoxia-induced radical injury to the walls of peripheral pulmonary arteries. In recent years, several studies have provided the solid evidence that radical tissue injury is a crucial mechanism in the development of HPH (for review see Hampl and Herget 2000). Release of both oxygen radicals (Hoshikawa *et al.* 1995, Nakanishi *et al.* 1995, Herget *et al.* 2000, Hoshikawa *et al.* 2001) and NO with its derivatives (Hampl and Herget 2000, Hampl *et al.* 2006) play a role. Production of NO and ROS peaks at the early phase of exposure to hypoxia (Hoshikawa *et al.*




Fig. 2. Four days of exposure to ambient air (N), hypoxia (H) and hypoxia with hypercapnia (H + CO₂). Data are means ± SEM, numbers of rats are in parentheses. **A.** amount of expired of NO of awake rats, **B.** concentration of NOx in plasma, **C.** concentration of plasma nitrotyrosine (marker of superoxide and NO interaction).




Fig. 3. Typical SDS-PAGE gel electrophoresis of collagen fraction isolated from peripheral pulmonary arteries of rats exposed to chronic hypoxia (H), normal control rat (C) and rats exposed to hypoxia and hypercapnia (H + CO₂). γ - gamma fraction, chain polymers of individual α chains of collagen type I; β - beta fraction, chain dimmers of individual α chains of collagen type I; α_1 - individual α_1 chains of collagen type I; α_2 - α_2 chains of collagen type I; **A1** - $\frac{3}{4}$ fragments of α_1 chains; **A2** - $\frac{3}{4}$ fragments of α_2 chains; **B1** - $\frac{1}{4}$ fragments of α_1 chains; **B2** - $\frac{1}{4}$ fragments of α_2 chains.

2001, Lachmanová *et al.* 2005, Hampl *et al.* 2006). Several groups reported a protective role of hypercapnia against free radical injury (Shibata *et al.* 1998, Laffey *et al.* 2000, Lavani *et al.* 2007, Skoumalová *et al.* 2008). The mechanism of this action may be linked to the stabilization of iron-transferin binding due to synergistic action of bicarbonate (Edeker *et al.* 1995) or to the interaction of CO₂ with peroxynitrite and the formation of nitrosoperoxycarbonate with a very short lifetime and therefore restricted diffusion area (Lyman and Hurst 1996, Uppu *et al.* 1996). We assume that peroxynitrite, a highly reactive product of superoxide-nitric oxide interaction (Huie and Padmaja 1993, Beckman and Koppenol 1996), triggers the collagen breakdown in hypoxia-induced injury of prealveolar pulmonary blood

vessels (Novotná and Herget 2002) and starts the process of remodeling. The finding that serum concentration of nitrotyrosine (marker of peroxynitrite formation (Beckman 1996) was significantly lower in the H + CO₂ group than in the hypoxic group without hypercapnia is, therefore, in agreement with our observation of less activated collagenolysis and less pronounced remodeling in vessels from rats exposed to hypoxia and hypercapnia.

The interpretation of our results is that hypercapnia inhibits hypoxia-induced tissue injury. Consequently, less NO and ROS is released and the vascular damage is less pronounced.

The alternative explanation is that HPH is decreased because of a vasorelaxant effect of CO₂. Belik *et al.* (2009) reported enhancement of endothelium-




Fig. 4. Relative density of collagen γ , β and α fraction and $\frac{3}{4}$ and $\frac{1}{4}$ fragments A1, A2, B1, B2 isolated from peripheral pulmonary arteries of rats exposed to chronic hypoxia (H), normal control rat (C) and rats exposed to hypoxia and hypercapnia (H + CO₂), separated by SDS-PAGE gel electrophoresis.

dependent vasodilatation by severe chronic hypercapnia (10 %). An increase in NO production was ascribed to downregulation of arginase expression and a subsequent increase in endothelial NO production (Belik *et al.* 2009). This alternative is not supported by our finding of less exhaled NO in the group exposed to hypoxia and hypercapnia than in the group kept in hypoxia alone. It should be noted, however, that our rats were exposed to much less severe hypercapnia (4 %).

In summary, we report that concomitant hypercapnia inhibits the development of chronic hypoxic pulmonary hypertension. This is in agreement with the generally accepted permissive effect of hypercapnia (Laffey and Kavanagh 1999).

References

- ASTIN TW: Airways obstruction and arterial blood gas tensions in chronic obstructive lung disease. *Respiration* **29**: 74-82, 1972.
- BECKMAN JS: Oxidative damage and tyrosine nitration from peroxynitrite. *Chem Res Toxicol* **9**: 836-844, 1996.
- BECKMAN JS, KOPPENOL WH: Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly. *Am J Physiol* **271**: C1424-C1437, 1996.
- BELIK J, STEVENS D, PAN J, SHEHNAZ D, IBRAHIM C, KANTORES C, IVANOVSKA J, GRASEMANN H, JANKOV RP: Chronic hypercapnia downregulates arginase expression and activity and increases pulmonary arterial smooth muscle relaxation in the newborn rat. *Am J Physiol* **297**: L777-L784, 2009.
- CANEPA A, CHAVEZ R, HURTADO A, ROTTA A, VELASQUEZ T: Pulmonary circulation at sea level and at high altitudes. *J Appl Physiol* **9**: 328-336, 1956.
- EDEKER BL, RASMUSSEN GT, BRITIGAN BE: Bicarbonate and phosphate ions protect transferrin from myeloperoxidase-mediated damage. *J Leukoc Biol* **58**: 59-64, 1995.
- EMERY CJ, SLOAN PJ, MOHAMMED FH, BARER GR: The action of hypercapnia during hypoxia on pulmonary vessels. *Bull Eur Physiopathol Respir* **13**: 763-776, 1977.
- HAMPL V, HERGET J: Perinatal hypoxia increases hypoxic pulmonary vasoconstriction in adult rats recovering from chronic exposure to hypoxia. *Am Rev Respir Dis* **142**: 619-624, 1990.

Conflict of Interest

There is no conflict of interest.

Acknowledgements

Supported by grants GACR 305/05/08/108 and GAUK 78007 as well as by Cardiovascular Research Center 1M510. Preliminary report of the part of results was presented at the Experimental Biology Meeting 2009, New Orleans, USA and Meeting of the Czech Physiological Society, 2001 and 2002, Prague, Czech Republic. We would like to thank to our lab assistants: Ms Olga Hniličková and Pavlína Smolková for assisting experiments and Ms Elizabeth "Liz" Coling help with the manuscript preparation.

- HAMPL V, HERGET J: Role of nitric oxide in the pathogenesis of chronic pulmonary hypertension. *Physiol Rev* **80**: 1337-1384, 2000.
- HAMPL V, BÍBOVÁ J, POVÝŠILOVÁ V, HERGET J: Dehydroepiandrosterone sulphate reduces chronic hypoxic pulmonary hypertension in rats. *Eur Respir J* **21**: 862-865, 2003.
- HAMPL V, BÍBOVÁ J, BAŇASOVÁ A, UHLÍK J, MIKOVÁ D, HNILIČKOVÁ O, LACHMANOVÁ V, HERGET J: Pulmonary vascular iNOS induction participates in the onset of chronic hypoxic pulmonary hypertension. *Am J Physiol* **290**: L11-L20, 2006.
- HERGET J, PALEČEK F: Pulmonary arterial blood pressure in closed chest rats. Changes after catecholamines, histamine and serotonin. *Arch Int Pharmacodyn Ther* **198**: 107-117, 1972.
- HERGET J, SUGGETT AJ, LEACH E, BARER GR: Resolution of pulmonary hypertension and other features induced by chronic hypoxia in rats during complete and intermittent normoxia. *Thorax* **33**: 468-473, 1978.
- HERGET J, WILHELM J, NOVOTNÁ J, ECKHARDT A, VYTÁŠEK R, MRÁZKOVÁ L, OŠTÁDAL M: A possible role of oxidant tissue injury in the development of pulmonary hypertension. *Physiol Res* **49**: 493-501, 2000.
- HERGET J, VYTÁŠEK R, BAŇASOVÁ A, BÍBOVÁ J, HAMPL V, FIŠÁRKOVÁ B: Hypercapnia inhibits hypoxic pulmonary hypertension in rats. *Physiol Res* **50**: P7, 2001.
- HERGET J, NOVOTNÁ J, BLÁHOVÁ L: Effect of hypercapnia on vascular remodelling induced by chronic hypoxia. *Physiol Res* **51**: 12P, 2002.
- HOSHIKAWA Y, ONO S, TANITA S, SAKUMA T, NODA M, TABATA T, UEDA S, ASHINO Y, FUJIMURA S: Contribution of oxidative stress to pulmonary hypertension induced by chronic hypoxia. *Nihon Kyobu Shikkan Gakkai Zasshi* **33**: 1169-1173, 1995.
- HOSHIKAWA Y, ONO S, SUZUKI S, TANITA T, CHIDA M, SONG C, NODA M, TABATA T, VOELKEL NF, FUJIMURA S: Generation of oxidative stress contributes to the development of pulmonary hypertension induced by hypoxia. *J Appl Physiol* **90**: 1299-1306, 2001.
- HOWELL K, OOI H, PRESTON R, MCLOUGHLIN P: Structural basis of hypoxic pulmonary hypertension: the modifying effect of chronic hypercapnia. *Exp Physiol* **89**: 66-72, 2004.
- HUIE RE, PADMAJA S: The reaction of NO with superoxide. *Free Radic Res Commun* **18**: 195-199, 1993.
- HULTGREN HN, GROVER RJ: Circulatory adaptation to high altitude. *Annu Rev Med* **16**: 119-152, 1968.
- KANTORES C, MCNAMARA PJ, TEIXEIRA L, ENGELBERTS D, MURTHY P, KAVANAGH BP, JANKOV RP: Therapeutic hypercapnia prevents chronic hypoxia-induced pulmonary hypertension in the newborn rat. *Am J Physiol* **291**: L912-L922, 2006.
- LAFHEY JG, KAVANAGH BP: Carbon dioxide and the critically ill – too little of a good thing? *Lancet* **354**: 1283-1286, 1999.
- LAFHEY JG, TANAKA M, ENGELBERTS D, LUO X, YUAN S, TANSWELL AK, POST M, LINDSAY T, KAVANAGH BP: Therapeutic hypercapnia reduces pulmonary and systemic injury following in vivo lung reperfusion. *Am J Respir Crit Care Med* **162**: 2287-2294, 2000.
- LACHMANOVÁ V, HNILIČKOVÁ O, POVÝŠILOVÁ V, HAMPL V, HERGET J: N-acetylcysteine inhibits hypoxic pulmonary hypertension most effectively in the initial phase of chronic hypoxia. *Life Sci* **77**: 175-182, 2005.
- LAPP M, SEATON A, KAPLAN L, MR H, MORGAN W: Pulmonary hemodynamics in symptomatic coal miners. *Am Rev Respir Dis* **104**: 418-429, 1971.
- LAVANI R, CHANG WT, ANDERSON T, SHAO ZH, WOJCIK KR, LI CQ, PIETROWSKI R, BEISER DG, IDRIS AH, HAMANN KJ, BECKER LB, VANDEN HOEK TL: Altering CO₂ during reperfusion of ischemic cardiomyocytes modifies mitochondrial oxidant injury. *Crit Care Med* **35**: 1709-1716, 2007.
- LYMAR SV, HURST JK: Carbon dioxide: physiological catalyst for peroxynitrite-mediated cellular damage or cellular protectant? *Chem Res Toxicol* **9**: 845-850, 1996.
- NAKANISHI I, TAJIMA F, NAKAMURA A, YAGURA SY, OOKAWARA T, YAMASHITA H, SUZUKI K, TANIGUCHI N, OHNO H: Effects of hypobaric hypoxia on antioxidant enzymes in rats. *J Physiol Lond* **489**: 869-876, 1995.
- NOVOTNÁ J, HERGET J: Exposure to chronic hypoxia induces qualitative changes of collagen in the walls of peripheral pulmonary arteries. *Life Sci* **62**: 1-12, 1998.

-
- NOVOTNÁ J, HERGET J: Possible role of matrix metalloproteinases in reconstruction of peripheral pulmonary arteries induced by hypoxia. *Physiol Res* **51**: 323-334, 2002.
- OOI H, CADOGAN E, SWEENEY M, HOWELL K, O'REGAN RG, McLOUGHLIN P: Chronic hypercapnia inhibits hypoxic pulmonary vascular remodeling. *Am J Physiol* **278**: H331-H338, 2000.
- PEPELKO WE: Effects of hypoxia and hypercapnia, singly and combined, on growing rats. *J Appl Physiol* **28**: 646-651, 1970.
- SHIBATA K, CREGG N, ENGELBERTS D, TAKEUCHI A, FEDORKO L, KAVANAGH BP: Hypercapnic acidosis may attenuate acute lung injury by inhibition of endogenous xanthine oxidase. *Am J Respir Crit Care Med* **158**: 1578-1584, 1998.
- SKOUMALOVÁ A, HERGET J, WILHELM J: Hypercapnia protects erythrocytes against free radical damage induced by hypoxia in exposed rats. *Cell Biochem Funct* **26**: 801-807, 2008.
- UPPU RM, SQUADRITO GL, PRYOR WA: Acceleration of peroxynitrite oxidations by carbon dioxide. *Arch Biochem Biophys* **327**: 335-343, 1996.
- WILHELM J, HERGET J: Hypoxia induces free radical damage to rat erythrocytes and spleen. Analysis of the fluorescent end-products of lipid peroxidation. *Int J Biochem Mol Biol* **31**: 671-681, 1999.
- WILKINSON M, LANGHORNE CA, HEATH D, BARER GR, HOWARD P: A pathophysiological study of 10 cases of hypoxic cor pulmonale. *Q J Med* **66**: 65-85, 1988.
-